

Sample Paper-02
Mathematics
Class – XII

Time allowed: 3 hours

Maximum Marks: 100

General Instructions:

- (i) All questions are compulsory.
- (ii) This question paper contains 29 questions.
- (iii) Question 1- 4 in Section A are very short-answer type questions carrying 1 mark each.
- (iv) Question 5-12 in Section B are short-answer type questions carrying 2 marks each.
- (v) Question 13-23 in Section C are long-answer-I type questions carrying 4 marks each.
- (vi) Question 24-29 in Section D are long-answer-II type questions carrying 6 marks each.

Section A

1. Give an example of a relation which is symmetric but not reflexive and transitive.
2. Is the measure of 5 seconds is scalar or vector?
3. What is the domain of $\sin^{-1} x$?
4. Show that $\begin{vmatrix} \sin 10^\circ & -\cos 10^\circ \\ \sin 80^\circ & \cos 80^\circ \end{vmatrix} = 1$

Section B

5. Find the principal value of $\cot^{-1}\left(-\frac{1}{\sqrt{3}}\right)$.
6. If $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ and $A^2 = I$. Find relation given by $\alpha^2 = I$.
7. Verify Rolle's Theorem for the function $f(x) = x^2 + 2x - 8, x \in [-4, 2]$
8. Find $\frac{dy}{dx} y = \tan^{-1}\left(\frac{\sin x}{1 + \cos x}\right)$
9. $\int \frac{\tan^4 \sqrt{x} \sec^2 \sqrt{x}}{\sqrt{x}} dx$
10. Verify that the function is a solution of the corresponding diff eq.
 $x + y = \tan^{-1} y; y^2 y' + y^2 + 1 = 0$
11. Find the angle between vectors \vec{a} and \vec{b} if $|\vec{a}| = \sqrt{3}, |\vec{b}| = 2, \vec{a} \cdot \vec{b} = \sqrt{6}$
12. A die thrown three times. Events A and B are defined as below.
A: 4 on the third throw
B: 6 on the first and 5 on the second throw.
Find the probability of A given that B has already occurred.

Section C

13. Solve: $3 \sin^{-1}\left(\frac{2x}{1+x^2}\right) - 4 \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) + 2 \tan^{-1}\left(\frac{2x}{1-x^2}\right) = \frac{\pi}{3}$
14. Differentiate $\sin^{-1}\left[\frac{2^{x+1}}{1+4^x}\right]$

15. Use differentiation to approximate $(25)^{\frac{1}{3}}$
16. Find a unit vector perpendicular to each of the vector $(\vec{a} - \vec{b})$ and $(\vec{a} + \vec{b})$ where $\vec{a} = i + j + k$ and $\vec{b} = i + 2j + 3k$.
17. If A and B are independent events such that $P(A \cup B) = 0.6$, $P(A) = 0.2$. Find $P(B)$
18. Probability of solving specific problem independently by A and B are $\frac{1}{2}$ and $\frac{1}{3}$ respectively of both try to solve the problem independently, find the probability that
(i) the problem is solved
(ii) Exactly one of them solves the problem.
19. Solve $\frac{dy}{dx} + \frac{2y}{3} = \frac{x}{\sqrt{y}}$
20. If $A = \begin{bmatrix} 0 & -\tan \alpha / 2 \\ \tan \alpha / 2 & 0 \end{bmatrix}$ and I is the identity matrix of order 2, show that
$$I + A = (I - A) \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$
21. If $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots}}}$ prove that $(2y - 1) \frac{dy}{dx} = \frac{1}{x}$.
22. For any two vectors \vec{a} and \vec{b} , prove that $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$.
23. Find the vector equation of the plane passing through the intersection of the planes $\vec{r} \cdot (2i + 2j - 3k) = 7$, $\vec{r} \cdot (2i + 5j + 3k) = 9$ and the point $(2, 1, 3)$.

Section D

24. Let $A = \mathbb{R} - \{3\}$ and $B = \mathbb{R} - \{1\}$. Consider the function $f : A \rightarrow B$ defined by $f(x) = \left(\frac{x-2}{x-3}\right)$. Is f one-one and onto? Justify your answer.
25. Integrate $\int (\sqrt{\cot x} + \sqrt{\tan x}) dx$
26. Find the shortest distance between the lines
$$\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k})$$
$$\vec{r} = (2\hat{i} - \hat{j} - \hat{k}) + \mu(2\hat{i} + \hat{j} + 2\hat{k})$$
27. Using integration find the area of the region $\{(x, y) : x^2 + y^2 \leq 2ax, y^2 \geq ax, x, y \geq 0\}$.
28. The sum of three numbers is 6. If we multiply the third number by 3 and add second number to it we get 11. By adding first and third numbers we get double of the second number. Represent the following information mathematically and solve using matrices.
29. A factory manufactures two types of machines A and B. Each type is made of certain metal. The factory has only 480kgs of this metal available in a day. To manufacture machine A, 10 kgs of metal is required and 20kgs required for B. Machine A and B require 15 and 10 minutes to be painted. Painting department can use only 400 minutes in a day. The factory earns profit of 10,500 on machine A and 9000 on machine B. State as a linear programming problem and maximizes the profit.